Observation of Z candidates in High Pt Muon Data Sample

Alexei Varganov, Dan Amidei, Ken Bloom, Eric James

University of Michigan
28 November 2001


Muon High Pt Data Sample

- Stream J Data taken before shutdown
- Data were processed by v 4.0.0 production code
- Skimmed by Christoph Paus using HighPtMuonStrip module
- The data files are stored on fcdfsgi2 in the area:
 - /cdf/data38b/s0/top/prod/strip/dst/
- The file names are *.Dst-MUON

Muon – Muon Selections

- Global Cuts
 - One and only one primary vertex in the event
 - Primary vertex Z position: |Z|<60 cm
- "Good Muon" cuts
 - Transverse momentum: pt > 12 GeV/c
 - Track impact parameter: |d|<0.2 mm with beam correction
 - Track delta z: |z-Z|<5cm
 - Number of COT axial hits > 25
 - Presence of CMU and stub matching: |dX| < 5 cm
 - Muon tower energy: em $< 2 \ GeV$, hadron $< 6 \ GeV$ em+hadron $> 0.1 \ GeV$
 - Isolation: coneR7Et=(Et in cone 0.7–Et of Muon) < 4 GeV
- Final cuts for every muon pair in the event
 - Event should pass Global cuts
 - One muon is good and the other passes isolation, pt, and charge sign cuts


25 muon – muon Z candidates



Muon – Track Selections

- Global Cuts
 - Minimum number of tracks: nTracks > 3
- "Good Muon" cuts
 - Transverse momentum: pt > 12 GeV/c
 - Track impact parameter: |d|<0.25 mm with beam correction
 - Number of COT axial hits > 20 and COT stereo hits > 20
 - Presence of CMU stub with track matching: |dX| < 5 cm
 - Muon tower energy: em $< 2 \; GeV$, hadron $< 6 \; GeV$ em+hadron $> 0.1 \; GeV$
 - Isolation: coneR7Et=(Et in cone 0.7–Et of Muon) < 4 GeV
- "Good Track" cuts are the same as "Good Muon" cuts but no muon stub presence is required
- Final cuts for every muon pair in the event
 - Event should pass Global cuts
 - Good track and muon with opposie charge sign and |zTrack-zMuon|<1.7 cm

60 muon – track Z candidates



Z candidate : Run = 123566 Event = 469584

